



GOOD PRACTICE MANAGEMENT

Floating Pennywort (*Hydrocotyle ranunculoides*)





GOOD PRACTICE MANAGEMENT GUIDE FOR Floating Pennywort (*Hydrocotyle ranunculoides*)


Other names: Water Pennywort

For ID guides and more information:

<http://www.nonnativespecies.org/factsheet/factsheet.cfm?speciesId=1766>

<https://www.cabi.org/isc/datasheet/28068>

http://www.qbank.eu/Plants/Controlsheets/Hydrocotyle_ranunculoides_office_guide.pdf



Floating Pennywort (*Hydrocotyle ranunculoides*)

Image : GB NNSS Crown Copyright 2009

Cover image: Snowdonia Park Authority

Version 2: May 2020



MANAGEMENT SUMMARY



Ecology and impact of Floating Pennywort

This plant can be difficult to detect in early stages of invasion, so often management only begins when plant is well established. The maximum growth period for Floating Pennywort is late summer/early autumn, when it forms extensive floating mats. This species overwinters on margins/banks as smaller/flatter plant. Floating Pennywort has proved difficult to control in the past due to its rapid growth rates, its ability to vegetatively regrow from seeds/roots and even a single node fragment, as well as its resistance to chemical control. Due to this high regeneration capacity of its shoots and fragments *H. ranunculoides* can spread downstream and disperse to new regions very easily by means of waterfowl via water courses and by human intervention. This species is difficult to completely eradicate from sites at anything larger than a local scale without a large amount of resources and it may be more practical to aim to control and prevent spread of Floating Pennywort.

Effective management: summary

A combination of physical and chemical treatment has seen some success in the past, particularly for localised population reduction. Cutting followed by glyphosate treatment was shown to be effective at controlling and removing Floating Pennywort. In 2013, Norfolk Non-Native Species Initiative (NNSI) has successfully eradicated Floating Pennywort from the River Waveney. This work involved both volunteers and contractors spraying and pulling Floating Pennywort in the river, and continual monitoring of sites to ensure the success of control work and monitor any grow-back.

Whatever treatment you choose, continuous monitoring should occur in the first year of treatment, followed by monitoring of any regrowth in the following spring and summer. Overwintering of untreated material should be avoided at all costs, as this results in very rapid spread within a catchment.

MANAGEMENT METHODS

Chemical

Method: Glyphosate @ 6 l/ha in 400 litres of water. Efficacy greatly increased with use of the adjuvant Codacide Oil @ 1 l/ha

Potential equipment requirements (excluding PPE): Knapsack sprayer, preferably with a long-lance.

Most suitable situation for method: Particularly useful for terrestrial growth, thus avoiding deoxygenation problems. Good for sites that have poor access for mechanical removal, or as a treatment following removal.

Efficacy: Good, but less effective in late summer when the rafts become denser. Late summer treatments are effective against regrowth, if the site was treated earlier in the year. Up to end of July is the best time to apply chemical control, but if it is necessary to minimise damage to non-target species, treatment can be delayed until end of growing season when other plants have died back. The efficacy of chemical treatment is greatly improved with the addition of an adjuvant such as Codacide Oil or Top Film.

Constraints: Requires AqHerb01 approval and NPTC PA1 & PA6 qualifications. Potential non-target damage. There is a risk of deoxygenation if large decomposing biomass is not removed

Timescale: Decomposition following treatment can take up to 6 weeks in slow flowing water. However, thick mats can act as barriers to smaller leaves and prevent control being fully carried out. Best practice management is to follow up initial treatment with spot treatment through a hand lance, or removal of live plants by hand 2-4 weeks after the first application of chemical treatment. One study ((Newman J.R. & Duenas M.A. 2010; Aldridge et al. 2017) suggests that glyphosate with Top Film is most effective up until July and that that glyphosate with Codacide Oil works better from July onwards).

When to manage Floating Pennywort with glyphosate

Jan	Feb	March	April	May	June	July	Aug	Sept	Oct	Nov	Dec

Mechanical

Areas where cutting/pulling takes place must be netted to ensure fragments don't spread downstream. All cut material must be removed from the water and safely disposed of, preferably on site, well away from water courses.

It is possible to strongly reduce Floating Pennywort with mechanical/physical methods in localised areas if treatment is repeated. However, physical methods have historically not been successful at completely eradicating this species. A good way to follow up mechanical cutting (to ensure removal success) is to go back and remove any remaining fragments by hand. This technique was successfully employed at Gillingham Marshes in Suffolk to significantly reduce cover of Floating Pennywort (Kelly 2006) using a mechanical digger for initial removal, then following up by hand-picking at least monthly throughout the growing season (March-September).

When follow up treatments are carried out, it is essential that nets/barriers used to collect plant fragments and any other equipment are cleaned thoroughly between treatments to avoid reinfection.

Cutting

Method: Weed-boat/reciprocating blades. Site netted to retain propagules.

Potential equipment requirements (excluding PPE): Specialist cutting equipment/weed boat. Vehicle & trailer if not disposing at site. Stop-nets and sweep nets.

Most suitable situation for method: Useful for reducing biomass of large infestations prior to chemical control.

Efficacy: Good, if propagules can be contained.

Constraints: Requires access if disposal is off-site. Expensive and it is likely this treatment would need to be repeated regularly, particularly in eutrophic situations. Avoid damage to the habitats of sensitive species, such as water voles and nesting birds.

Timescale: Cutting between July – Oct stops species becoming dominant.

Jan	Feb	March	April	May	June	July	Aug	Sept	Oct	Nov	Dec

Mechanical (cont)

Mechanical pulling

Method: Pulling using a hydraulic rake or bucket. Location netted to retain propagules

Potential equipment requirements (excluding PPE): weed-boat/tractor fitted with hydraulic rake or bucket. Vehicle & trailer if not disposing at site. Stop-nets and sweep nets.

Most suitable situation for method: Useful for reducing biomass of large infestations, prior to chemical control or hand pulling. Has better long-term control on established infestations than cutting.

Efficacy: Good, particularly when the rafts are mature

Constraints: Requires access. Avoid damage to the habitats of sensitive species, such as water voles and nesting birds. Can be expensive.

Timescale: Pull during growing season (March-Oct)

Jan	Feb	March	April	May	June	July	Aug	Sept	Oct	Nov	Dec

Hand pulling

Method: Hand-pulling, disposing of material by composting away from water habitats.

Potential equipment requirements (excluding PPE): Boats, drysuits, wheelbarrows, forks, rakes. Vehicle & trailer if not disposing at site. Stop-nets and sweep nets.

Most suitable situation for method: Particularly relevant for smaller infestations, but can be very effective against any size of infestation, depending on the resources and time available. Sites with good access or sensitive areas where non-target damage is an issue. Suitable for volunteer groups.



Mechanical (cont)

Hand pulling (cont)

Efficacy: Good, if propagules can be contained and resources exist to regularly repeat the task.

Constraints: Time-consuming (must be done carefully to ensure fragments do not break off), and requires good access

Timescale: Can be done throughout the year

Jan	Feb	March	April	May	June	July	Aug	Sept	Oct	Nov	Dec

Disposal

The Regulatory Statement with guidelines for treatment and disposal of invasive non-native plants is available at:

<https://www.gov.uk/government/publications/treatment-and-disposal-of-invasive-non-native-plants-rps-178/treatment-and-disposal-of-invasive-non-native-plants-rps-178>

Biocontrol

CABI is currently researching potential methods of biocontrol of Floating Pennywort with 2 control agents (the stem mining fly, *Eugaurax floridensis*, and the weevil *Listronotus elongatus*), but this is not widely available yet and the efficacy of these agents is still not certain. Research is ongoing and a test release may happen soon.



Preventing spread

Once escaped in the wild, fragmentation and dispersal of Floating Pennywort, (*Hydrocotyle ranunculoides*) can result from a variety of management and recreational activities. As this species is difficult to eradicate, raising awareness and practicing good biosecurity is key to effective management and preventing further spread. For guidance on this, please follow the Check, Clean, Dry guidance in the link below:

<http://www.nonnativespecies.org/checkcleandry/index.cfm>

Ineffective control

From the current evidence, the following methods are widely considered to be ineffective or untested at controlling Floating Pennywort:

Shading - not a viable option for control

Increasing flow of water - may restrict the growth of Floating Pennywort *in situ*, but this is not recommended as it may increase the spread of the plant downstream

Grazing - animals tend to only remove the leaves and not the whole plant, thus not solving the problem of vegetative fragments

Dredging - does not seem to be an effective method

Glyphosate used without adjuvant - has been shown in trials to have a negligible effect on controlling Floating Pennywort. Furthermore, using glyphosate without an approved adjuvant is not recommended as the herbicide is also rapidly excreted from the plant.

Hydrogen peroxide - has some visual encouraging effects, but does not in fact kill the plants.



Ineffective Control cont.

2,4 D amine – this chemical has been used for controlling Floating Pennywort in the past, but is not recommended due to difficulties caused by rapid translocation and excretion of the herbicide by the plant.

Increasing salinity, liquid nitrogen treatment and using flame treatments — thought to have some effects on reducing Floating Pennywort, but as these have not been widely tested and are not recommended.

Legislation

Floating Pennywort feather is listed under Schedule 9 to the Wildlife and Countryside Act 1981 with respect to England, Wales and Scotland. It is covered by Article 15 (2) of the Wildlife (Northern Ireland) Order 1985 as well. As such, it is an offence to plant or otherwise cause this species to grow in the wild. This species is also banned from sale in England due to its significant negative impacts on biodiversity and the economy. This is covered under [The Wildlife and Countryside Act 1981 \(prohibition on Sale etc. of Invasive Non-native Plants\) \(England\) Order 2014](#)

Floating pennywort is also on the EU List of Invasive Alien Species of Union Concern. The EU [Regulation \(1143/2014\) on invasive alien \(non-native\) species](#). Under the EU Invasive Alien Species Regulation it is also an offence to import into the EU, keep, grow or cultivate, transport (to, from or within the EU; except to facilitate eradication), place on the market, use or exchange this species - unless there are specific exemption or permit

A link to other resources on legislation of INNS:

<http://www.nonnativespecies.org/index.cfm?sectionid=23>



Health and Safety

Use of glyphosate requires AqHerb01 approval and NPTC PA1 & PA6 qualifications.

[Application to use herbicides in or near water](#)

[City & Guilds Level 2 Principles of Safe Handling and Application of Pesticides \(PA1\)](#)

[City & Guilds Level 2 Award in the Safe Application of Pesticides using Pedestrian Hand Held Equipment](#)

[Health and Safety Executive Code of Practice for Plant Protection Products](#)

Useful resources and guidance on health and safety when planning a project working with invasive species is available on the GBNNSS website:

<http://www.nonnativespecies.org/index.cfm?pageid=266>



References

Aldridge, D.C., Aldridge, S.L., Mead, A., Ockendon, N., Rocha, R., Scales, H., Smith, R.K., Zieritz, A. & Sutherland, W.J. (2017) Control of freshwater invasive species: global evidence for the effects of selected interventions. Synopses of Conservation Evidence Series. University of Cambridge, Cambridge, UK. <https://doi.org/10.11647/OBP.0131>

Brook Lyndhurst (2015) Local Action Groups for Managing Invasive Non-Native Species: A research review for Defra. www.nonnativespecies.org/downloadDocument.cfm?id=1417

CABI (2017) Progress with Weed Biocontrol projects CABI in the UK. https://www.cabi.org/Uploads/CABI/projects/WFD_update_May2017.pdf

Environment Agency (2010) Managing invasive non-native plants in or near fresh water. www.nonnativespecies.org/downloadDocument.cfm?id=1010

GT IBMA (2015) Invasive alien species in aquatic environments. In *Practical information and management insights, Vol. 2*. Emmanuelle Sarat, Emilie Mazaubert, Alain Dutartre, Nicolas Poulet and Yohann Soubeyran (editors) . <https://pipap.sprep.org/content/invasive-alien-species-aquatic-environments-practical-information-and-management-insights>

Hussner, A., Denys, L. and van Valkenburg, J. (2012): NOBANIS – Invasive Alien Species Fact Sheet – *Hydrocotyle ranunculoides*. Online Database of the European Network on Invasive Alien Species. www.nobanis.org

Hydrocotyle ranunculoides L. f. A guide to Identification, Risk Assessment and Management Plant Protection. (2011) Plant Protection Service, Wageningen, NL and Centre for Ecology and Hydrology, Wallingford, UK. http://www.q-bank.eu/Plants/Controlsheets/Hydrocotyle_ranunculoides_office_guide.pdf

Joost van der Burg, W. & Michielsen, J.M. (2010) Effect of flaming on *Hydrocotyle ranunculoides* L.f. survival. *Plant Research International*, Wageningen. 12pp.



Kelly, A., 2006. Removal of invasive Floating Pennywort *Hydrocotyle ranunculoides* from Gillingham Marshes, Suffolk, England. *Conservation Evidence*, 3, pp.52-53.

Newman J.R. & Duenas M.A. (2010) Information sheet: control of Floating Pennywort (*Hydrocotyle ranunculoides*). Centre for Ecology & Hydrology, Wallingford, UK. 3pp.

Scottish Natural Heritage (2014) Integrated Pest Management in Nature Conservation Handbook. <https://www.nature.scot/integrated-pest-management-nature-conservation-handbook>

Acknowledgements

Thanks to the GB NNS and The Environment Agency for input and advice.



Where To Go For More Information

- ◆ <http://www.anglingtrust.net/>
- ◆ <https://www.cabi.org/>
- ◆ <http://www.invasive-species.org/>
- ◆ <http://www.europe-aliens.org/>
- ◆ <http://www.nonnativespecies.org/beplantwise>
- ◆ <http://www.nonnativespecies.org/home>

RAPID

RAPID is a three year EU funded LIFE project led by the Animal and Plant Health Agency (APHA), with Natural England and Bristol Zoological Society as key partners that piloting innovative approaches to Invasive Alien Species (IAS) management in freshwater aquatic, riparian and coastal environments across England. The project is supported by a number of further Technical Partners.

<http://www.nonnativespecies.org/rapid>